

# **Lecture**

# **Module 8: Nutrient Cycling**

**Stan Gregory**  
**Department of Fisheries and Wildlife**  
**Oregon State University**

## **Learning Objectives**

Upon completion of this module, the participant will be able to:

1. Describe the various biotic and abiotic pathways of the major nutrients (C, N, P, K) through terrestrial and aquatic ecosystems.
2. Describe oligotrophic and eutrophic aquatic ecosystems and the processes that lead to these conditions.
3. Understand the spatial scale of effects of nutrient enrichment and depletion on biotic communities and their social and economic consequences.

## **Lecture Outline**

Major biologically important nutrients

Carbon

Sources

Dissolved vs. particular organic carbon (DOC vs POC)

C/N ratios

Refractory vs. labile compounds

Nitrogen

Nitrogen cycle

Phosphorus

Sources

Redfield ratio (N/P)

Nutrient cycling

Nutrient spiraling

Flux and uptake

Retention

Hypoxia in the Gulf of Mexico

Causes and consequences

Economic impact

Sources of nitrogen inputs—Upper Midwest

Changes in Mississippi

Management implications and solutions

## Exercises

### A) Classroom Exercises:


1. A point-source discharge increases the concentration of nitrate to 1 mg/L of NO<sub>3</sub>-N. You go downstream 100 m and find that the concentration is 0.70 mg/L. What is the average travel distance of a nitrate molecule in this stream?
2. For each of the types of nitrogen transformation described below, fill in the missing information. “Transformation” refers to the name of the process. “Source” refers to the initial form of nitrogen and “product” refers to the final form of nitrogen for that type of transformation. “Organism” refers to the type of organism that is capable of each type of transformation (be as specific as possible). For the type of organism indicated, indicate whether the process occurs in aerobic habitats, anaerobic habitats, or both.

Transformation	Source	Product	Organism	Aerobic or Anaerobic
	NO <sub>3</sub> -	N <sub>2</sub>	Bacteria	
	N <sub>2</sub>			Aerobic
	Dissolved organic N	NH <sub>4</sub> +	Bacteria	Both
			Nitrobacter	

### B) Field Exercise: Retention in Streams

The interface between terrestrial and aquatic ecosystems—riparian areas—strongly influences the production and input of food resources. Terrestrial ecosystems deliver large amounts of allochthonous organic material to streams and rivers. While the forms and amounts of allochthonous inputs are critical components of the food base, this material must be retained before it can be consumed or used as physical habitat. The process of retention is the removal of either dissolved or particulate material from transport by physical trapping of particulate matter, deposition of particulate matter, biological uptake of both dissolved and particulate material, or physical adsorption of dissolved material. Complex, rough stream channels retain material in transport more efficiently than simple, uniform channels.

After a quantity of material enters a stream, the proportion suspended in the water column will decrease exponentially as it is transported downstream (Figure 1).



**Figure 1.** Percent of leaves in transport as a function of distance downstream from the point of entry.

The retention of material in transport in a stream is based on a simple negative exponential model. This same relationship also is used to represent light extinction, decomposition of detritus, biotic uptake of dissolved nutrients, and radioactive decay (Equation 1).

(Equation 1) 
$$L_d = L_0 e^{-kd}$$

or

$$\ln L_d = \ln L_0 - kd$$

where  $L_d$  = number of leaves in transport at distance  $d$   
 $L_0$  = original number of leaves in transport  
 $k$  = instantaneous retention coefficient  
 $d$  = distance

Based on this negative exponential relationship, a constant proportion of the leaves will be removed from transport on a given increment of distance downstream. The retention coefficient,  $k$ , is the instantaneous rate of retention. The average travel distance for a particle can be calculated by the inverse of the retention coefficient,  $1/k$ .

## Procedure

### Leaf retention

1. Each group will be assigned to a specific 45-m section within the reach and will place a seine at the bottom of the 45-m section.
2. Leaves will be counted before the lab and soaked in water for 24 hr to make certain that all leaves are neutrally buoyant. Ginkgo leaves (*Ginkgo biloba*) will be used as tracer leaves because they are not commonly found along most streams, they are bright yellow and can be seen easily, and they are tough and resistant to fragmentation when dried and stored.
3. The leaves will be released into the stream carefully distributing them across the channel and avoiding clumping.



### Dye

1. After 1 hr, all leaves that reached the bottom net will be collected and counted.
2. Immediately after the initial leaf release, each group will dissolve 5 g of fluorescein dye in 1 L of stream water.
3. A person at the top end of the section will coordinate with a person at the lower end. At a predetermined signal the dye will be poured into the center of the channel at the top of the section.
4. The person at the lower end will record the time that it took the dye to appear visually at the lower end of the section as a measure of maximum velocity.
5. They will also record the time at which the dye is flushed from the section and can no longer be seen. The time that is halfway between the maximum velocity and final clearance will be used as a measure of median velocity.

### Measures to be Included in Project Reports

Calculate the retention coefficients and travel distances for the four 45-m sections and determine averages for the larger study reach.

- Retention coefficient for each section (45 m)
- Average leaf travel distance for each section (45 m)
- Average retention coefficients for the reach
- Average leaf travel distances for the reach
- Maximum velocity for each section
- Median velocity for each section
- Average maximum velocity for the reach
- Average median velocity for the reach

## **Study Questions**

1. Nutrients such as carbon, nitrogen, and phosphorus undergo intricate cycles in terrestrial systems, all of which are influenced by soils, plants, water, bacteria and animals. What attributes of terrestrial systems are important for efficient cycling of nutrients?
2. Define the concept of nutrient spiraling. Provide a general equation to explain your definition.




## References and Selected Reading

- Alexander, R. B., R. A. Smith, and G. E. Schwarz. 2000. Effect of stream channel size on the delivery of nitrogen to the Gulf of Mexico. *Nature* 403:758-761.
- Binkley, D. and T. C. Brown. 1993. Forest practices as nonpoint sources of pollution in North America. *Water Resources Bulletin* 29:729-740.
- Duff, J. H., and F. J. Triska. 1990. Denitrification in sediments from the hyporheic zone adjacent to a small forested stream. *Canadian Journal of Fisheries and Aquatic Sciences* 47:1140-1147.
- Grimm, N. B., and S. G. Fisher. 1984. Exchange between interstitial and surface water: Implications for stream metabolism and nutrient cycling. *Hydrobiologia* 111:219-228.
- Groffman, P. M. and J. M. Tiedje. 1989. Denitrification in north temperate forest soils: spatial and temporal patterns at the landscape and seasonal scales. *Soil Biological Biochemistry* 21(5):613-620.
- Hill, A. R. 1983. Denitrification: its importance in a river draining an intensively cropped watershed. *Agriculture, Ecosystems, and Environment* 10:47-62.
- Hill, A. R. 1996. Nitrate removal in stream riparian zones. *J. Environ. Qual.* 25:743-755.
- Lowrance, R. 1992. Groundwater nitrate and denitrification in a coastal plain riparian forest. *Journal of Environmental Quality* 21:401-405.
- Peterjohn, W. T., and D. L. Correll. 1984. Nutrient dynamics in an agricultural watershed: observations on the role of a riparian forest. *Ecology* 65:1466-1475.
- Pinay, G., A. Fabre, P. Vervier, and F. Gazelle. 1992. Control of carbon, nitrogen, phosphorous distribution in soils of riparian forests. *Landscape Ecology* 6:121-132.
- Schnabel, R. R., L. F. Cornish, W. L. Stout and J. A. Shaffer. 1996. Denitrification in a grassed and a wooded, valley and ridge, riparian ecotone. *Journal of Environmental Quality* 25:1230-1235.
- Showstack, R. 2000. Nutrient over-enrichment implicated in multiple problems in U.S. waterways. *Eos, Transactions, American Geophysical Union* 81(43):497-499.
- Vitousek P. M. , J. D. Aber, R. W. Howarth, G. E. Likens, P. A. Matson, D. W. Schindler, W. H. Schlesinger, and D. G. Tilman. 1997. Human alteration of the global nitrogen cycle: Sources and consequences. *Ecol. Appl.* 7:737-750.



Slides used in lecture

Slide  
1



Slide 1: Major Nutrients

- Carbon
- Nitrogen
- Phosphorus

Understanding the Landscape

---

---

---


---

---

---

---

Slide  
2



Slide 2: Organic Carbon

Sources:

- Plant production within the ecosystem
- Plant production outside the ecosystem

Understanding the Landscape

---

---

---


---

---

---

---

Slide  
3



Slide 3: Organic matter in aquatic ecosystems

- Dissolved organic matter  
< 0.45  $\mu\text{m}$
- Particulate organic matter  
> 0.45  $\mu\text{m}$

Understanding the Landscape

---

---

---

---

---

---

---



Slide  
4

### Comparison of DOC and POC

In aquatic ecosystems, the amount of DOC usually greatly exceeds that of POC:

DOC/POC ~ 5 in oligotrophic lakes  
DOC/POC ~ 10 in eutrophic lakes

Understanding the Landscape

---

---

---

---

---

---

---

Slide  
5

### Carbon to Nitrogen Ratios

≈ Index of quality of organic matter

**Terrestrial organic matter:**  
high C/N ratios (20-200)

**Aquatic plants:**  
low C/N ratios (5-15)

Understanding the Landscape

---

---

---

---

---

---

---

Slide  
6

### Composition of Organic Matter

**Refractory Compounds:**

- Resist decay
- High molecular weight
- Low solubility
- Largely structural

Understanding the Landscape

---

---

---

---


---

---

---



Slide  
7




---

---

---

---

---

---

---

---

Slide  
8

### Composition of Organic Matter

**Labile Compounds:**

- Utilized rapidly
- Low molecular weight
- High solubility
- Non-structural

Understanding the Landscape

---

---

---

---


---

---

---

---

Slide  
9




---

---

---

---

---

---

---

---



Slide 10

Dissolved Organic Carbon

- Labile compounds are used rapidly
- Concentrations tend to increase downstream
- Slight seasonal fluctuation

Understanding the Landscape

---

---

---

---

---

---

---

Slide 11

Nitrogen

- Nitrogen gas -  $N_2$
- Ammonia -  $NH_3$
- Ammonium -  $NH_4^+$
- Nitrite -  $NO_2^-$
- Nitrate -  $NO_3^-$
- Organic N

Understanding the Landscape

---

---

---


---

---

---

---

Slide 12



---

---

---

---


---

---

---



Slide  
13



## Phosphorus

- Inorganic phosphorus
  - Orthophosphate
  - Soluble reactive phosphorus
- Organic phosphorus

Understanding the Landscape

---

---

---


---

---

---

---

Slide  
14



## Sources for aquatic ecosystems

- Groundwater
  - Concentrations are generally low
  - Often less than 20 ug/l
- Surface runoff
  - Soil related
  - Often a major source of input
  - Related to soils, topography, vegetation cover, quantity of runoff, land use, and pollution

Understanding the Landscape

---

---

---


---

---

---

---

Slide  
15



## Sources for lakes

- Phosphorus content of parent geology is highly variable
- Pollution from fertilizers is a major source of phosphorus
- Pollution from detergents is a major source of phosphorus

Understanding the Landscape

---

---

---

---

---

---

---



Slide 16

Sediments

- Concentrations of phosphorus often increase near sediments because of the effects of reduced oxygen concentration on redox potential
- Areas around habitats with low oxygen often contain higher phosphorus concentrations

Understanding the Landscape

---

---

---

---

---

---

---

Slide 17

Redfield Ratio

- Determined by ratio of N and P in cells
- N/P molar ratio
- < 15 photosynthesis is N limited
- > 30 photosynthesis is P limited

Understanding the Landscape

---

---

---

---

---

---

---

Slide 18

Nutrient Cycling

S = Flux/Uptake

$S = N \cdot v / N \cdot u$

N = Standing stock (g/m)  
v = Velocity (m/s)  
u = Uptake rate (s<sup>-1</sup>)  
S = m

Understanding the Landscape

---

---

---

---


---

---

---



Slide  
19




---

---

---

---

---

---

---

---

Slide  
20

**Retention**

$$M_d = M_o * e^{-kd}$$

- $M_d$  = Mass at distance
- $M_o$  = Mass at original location
- $e$  = Natural logarithm
- $k$  = Instantaneous retention constant
- $d$  = Distance downstream

---

---

---

---

---

---

---

---

Slide  
21

**Retention**

- Average travel distance =  $1/k$
- Average travel distance  $\approx S_w$

---

---

---

---

---

---

---

---



Slide  
22



---

---

---

---

---

---

---

Slide  
23



---

---

---


---

---

---

---

Slide  
24



---

---

---

---


---

---

---



Slide  
25




---

---

---

---

---

---

---

---

Slide  
26

### Hypoxia in Gulf of Mexico

- Zone was 20,000 km<sup>2</sup> (7,728 mi<sup>2</sup>) in 1999
- Size of New Jersey
- Zone was 4,000 km<sup>2</sup> (1,545 mi<sup>2</sup>) in 2000
- Extended to depths of 100 ft

Understanding the Landscape

---

---

---

---

---

---

---

---

Slide  
27

### Economic Implications of Hypoxia Gulf of Mexico

- Spawning grounds, migration pathways, feeding habitat, etc.
- \$2.8 billion fishery

---

---

---

---

---


---

---

---



Slide  
28



---

---

---

---

---

---

---

Slide  
29



---

---

---


---

---

---

---

Slide  
30



---

---

---

---


---

---

---



Slide  
31




---

---

---

---


---

---

---

---

Slide  
32




---

---

---

---


---

---

---

---

Slide  
33




---

---

---

---

---


---

---

---



Slide  
34



---

---

---

---

---

---

---

Slide  
35



---

---

---

---

---

---

---

Slide  
36



---

---

---

---


---

---

---



Slide  
37



---

---

---

---

---

---

---

Slide  
38



---

---

---

---

---

---

---

Slide  
39



---

---

---

---

---

---

---



Slide  
40



---

---

---


---

---

---

---

Slide  
41



---

---

---


---

---

---

---

Slide  
42



---

---

---

---


---

---

---



Slide  
43



## Solutions?

### On-Farm Practices

- Application rate  
("insurance applications")
- Residual fertilizer, legume input
- Fertilizer content & applications for manure

**Only 15-20% reduction**

---

---

---

---

---

---

---

---

Slide  
44



## Solutions?

### Wetland & Riparian Restoration

One of the most effective nutrient barriers in agricultural regions

- 5 million acres of wetland restoration
- 19 million acres of riparian restoration

**Additional 40% reduction**

---

---

---

---


---

---

---

---

Slide  
45




---

---

---

---

---


---

---

---



Slide  
46



---

---

---

---

---

---

---